




AP Calculus BC – AP Exam Review Project –Free Response Problems → Group 5

Year & #	2017 2	2015 3	2013 4	2011 3	2010 5	
National Average	3.21	5.71	4.62	1.67	2.44	
Type of Problem	Polar	Bicycle Integral	Max Min from Graph	Perimeter & Volume	Euler's Method	Taylor Twice Differential



The figure above shows the polar curves $r = f(\theta) = 1 + \sin \theta \cos(2\theta)$ and $r = g(\theta) = 2 \cos \theta$ for $0 \leq \theta \leq \frac{\pi}{2}$. Let R be the region in the first quadrant bounded by the curve $r = f(\theta)$ and the x -axis. Let S be the region in the first quadrant bounded by the curve $r = f(\theta)$, the curve $r = g(\theta)$, and the x -axis.

- Find the area of R .
- The ray $\theta = k$, where $0 < k < \frac{\pi}{2}$, divides S into two regions of equal area. Write, but do not solve, an equation involving one or more integrals whose solution gives the value of k .
- For each θ , $0 \leq \theta \leq \frac{\pi}{2}$, let $w(\theta)$ be the distance between the points with polar coordinates $(f(\theta), \theta)$ and $(g(\theta), \theta)$. Write an expression for $w(\theta)$. Find w_A , the average value of $w(\theta)$ over the interval $0 \leq \theta \leq \frac{\pi}{2}$.
- Using the information from part (c), find the value of θ for which $w(\theta) = w_A$. Is the function $w(\theta)$ increasing or decreasing at that value of θ ? Give a reason for your answer.

t (minutes)	0	12	20	24	40
$v(t)$ (meters per minute)	0	200	240	-220	150


Johanna jogs along a straight path. For $0 \leq t \leq 40$, Johanna's velocity is given by a differentiable function v . Selected values of $v(t)$, where t is measured in minutes and $v(t)$ is measured in meters per minute, are given in the table above.

- (a) Use the data in the table to estimate the value of $v'(16)$.
- (b) Using correct units, explain the meaning of the definite integral $\int_0^{40} |v(t)| dt$ in the context of the problem.

Approximate the value of $\int_0^{40} |v(t)| dt$ using a right Riemann sum with the four subintervals indicated in the table.


- (c) Bob is riding his bicycle along the same path. For $0 \leq t \leq 10$, Bob's velocity is modeled by $B(t) = t^3 - 6t^2 + 300$, where t is measured in minutes and $B(t)$ is measured in meters per minute. Find Bob's acceleration at time $t = 5$.
- (d) Based on the model B from part (c), find Bob's average velocity during the interval $0 \leq t \leq 10$.

The figure above shows the graph of f' , the derivative of a twice-differentiable function f , on the closed interval $0 \leq x \leq 8$. The graph of f' has horizontal tangent lines at $x = 1$, $x = 3$, and $x = 5$. The areas of the regions between the graph of f' and the x -axis are labeled in the figure. The function f is defined for all real numbers and satisfies $f(8) = 4$.



- (a) Find all values of x on the open interval $0 < x < 8$ for which the function f has a local minimum. Justify your answer.
- (b) Determine the absolute minimum value of f on the closed interval $0 \leq x \leq 8$. Justify your answer.
- (c) On what open intervals contained in $0 < x < 8$ is the graph of f both concave down and increasing? Explain your reasoning.
- (d) The function g is defined by $g(x) = (f(x))^3$. If $f(3) = -\frac{5}{2}$, find the slope of the line tangent to the graph of g at $x = 3$.

Let $f(x) = e^{2x}$. Let R be the region in the first quadrant bounded by the graph of f , the coordinate axes, and the vertical line $x = k$, where $k > 0$. The region R is shown in the figure above.



- Write, but do not evaluate, an expression involving an integral that gives the perimeter of R in terms of k .
- The region R is rotated about the x -axis to form a solid. Find the volume, V , of the solid in terms of k .

- The volume V , found in part (b), changes as k changes. If $\frac{dk}{dt} = \frac{1}{3}$, determine $\frac{dV}{dt}$ when $k = \frac{1}{2}$.

Consider the differential equation $\frac{dy}{dx} = 1 - y$. Let $y = f(x)$ be the particular solution to this differential equation with the initial condition $f(1) = 0$. For this particular solution, $f(x) < 1$ for all values of x .

- Use Euler's method, starting at $x = 1$ with two steps of equal size, to approximate $f(0)$. Show the work that leads to your answer.
- Find $\lim_{x \rightarrow 1} \frac{f(x)}{x^3 - 1}$. Show the work that leads to your answer.
- Find the particular solution $y = f(x)$ to the differential equation $\frac{dy}{dx} = 1 - y$ with the initial condition $f(1) = 0$.